

Kaeriyama, H., Asahi, T., Nakaoka, M., Ichimi, K., Hashimoto, T., Yamada, T. and Tada, K. 2010.

Size Distribution of the Phytoplankton Community and Standing Stock of Bacteria in Hiroshima Bay during the Autumn Season.

Bulletin on Coastal Oceanography 48(1): 65-74

Komai, T. 2010.

A Review of the Northwestern Pacific Species of the genus *Paguristes* (Decapoda: Anomura: Diogenidae). III. Clarification of the Identity of a Species Heretofore Referred to *Paguristes balanophilus* Alcock and Descriptions of Two New Species from Japan.

Nat. Hist. Res. 11(1): 09-33

Mizuno, K., Shoji, J., Zenitani, H., Kono, N., Maehara, T. and Kishida, T. 2010.

Spatial distribution of Japanese sardine, gizzard shad and Japanese anchovy larvae in the Sea of Hiuchi, central Seto Inland Sea, in the 1990s.

Bull. Jpn. Soc. Fish. Oceanogr. 74(3): 159-166

Nakajima, S., Kanayama, S., Sekimoto, T. and Iseki, K. 2010.

A study on the formation mechanism of isolated cold dome and physical environment near the bottom of Suo-Nada in summer.

Journal of Japan Society of Civil Engineers, Ser. B2 66(1): 951-955

Nakanishi, T., Iseki, K., Miyashita, Y., Koike, K., Hamaguchi, M. and Tezuka, N. 2010.

Summer distribution and short-term variation of the bottom turbid layer in Suo-sound in the Western Seto Inland Sea, Japan.

Biosphere Sci. 49: 21-30

Olasehinde, E. F., Takeda K. and Sakugawa, H. 2010.

Photochemical Production and Consumption Mechanisms of Nitric Oxide in Seawater.

Environ. Sci. Technol. 44: 8403-8408

Saito, H. and v. Salvini-Plawen, L. 2010.

A New Species of *Anamenia* (Mollusca: Solenogastres: Cavibelonia) from Southern Japan.

VENUS 69 (1-2): 1-15

Sakai, Y., Ochi, Y., Tsuboi, M., Kadota, K., Shimizu, N., Shoji, J., Matsumoto, K., Mabuchi, K., Kuniyoshi, H., Ohtsuka, S. and Hashimoto, H. 2010.

Fish fauna of shallow waters of Aki Nada, Seto Inland sea, Japan.

Biosphere Sci. 49: 7-20

- Takeda, M. and Marumura, M. 2010.
Spider Crabs of the Genus *Huenia* De Haan, 1837 (Crustacea, Decapoda, Brachyura) from Japan, with Descriptions of Two New Species.
Bull. Natl. Mus. Nat. Sci., Ser. A 36(2): 39–48
- Tanaka, G. and Seto, K. 2010.
Description of appendages from three marine trachyleberidid species (Ostracoda, Crustacea) from Japan.
Journal of Micropalaeontology, 29: 5-16
- Tomikawa, K., Kakui, K. and Yamasaki, H. 2010.
A New Species of *Psammogammarus* (Amphipoda: Melitidae) from Kuchinoerabu Island, Japan, with a Note on Its Feeding Habits.
Zoological Science 27(7): 615-626
- Ueoka, R., Ise, Y., Ohtsuka, S., Okada, S., Yamori, T. and Matsunaga, S. 2010.
Yaku'amides A and B, Cytotoxic Linear Peptides Rich in Dehydroamino Acids from the Marine Sponge *Ceratopsion* sp.
J. AM. CHEM. SOC. 132: 17692–17694
- Uye, S. 2010.
Studies on Functional Roles of Zooplankton in Coastal Marine Ecosystem: Toward Restoring Productive Seas for Global Sustainability.
Oceanography in Japan 19(6): 283– 299
- Yang, CH and Chan, TY 2010.
A new slipper lobster of the Genus *Galearctus* Holthuis, 2002 (Decapoda: Scyllaridae) from Taiwan and Japan.
Studies on Malacostraca: Lipke Bijdeley Holthuis Memorial Volume: 735-745